

Evaluation of correlation-ship between testicular volumes and semen parameters among healthy adults

Arunkumar Neelakandan^{1*}, Alex Daniel Prabhu Arul Pitchai², Einstein Arulraj³

^{1,2,3}Department of Radiology, Chettinad Hospital & Research Institute, Kelambakkam, Chennai, Tamil Nadu, INDIA.

Email: dr.arun.n@gmail.com

Abstract

Background: Assessment of testicular size and semen parameters are an essential and initial method for the evaluation of testicular function. For accurate measurement of testicular volume, ultrasonography is considered as most crucial radiological technique. Our study aims to determine variations in testicular volumes among healthy adults with the help of ultrasonography. **Materials and Method:** Our study was a prospective observational category study. It was carried out at the Department of Radiology, Chettinad Hospital and Research Institute, Kelambakkam, Chennai. This study was carried between August 2019 and August 2021. This study was carried between August 2019 and August 2021. A total of 100 adult males aged 18 to 70 years were assessed to measure the testicular size, and Lambert's formula was used to calculate testicular volumes on both sides. All study participants voluntarily underwent serum hormone analysis, conventional semen parameters evaluation. The relationship between testicular volumes, semen parameters, anthropometric measurements and other characteristics of participants were also evaluated. **Results:** Lambert's formula was used, and the testicular volumes were determined. A total number of 100 adult males aged 18 to 70 years (median age 38 years) participated in this study. The mean testicular volume was recorded as $21.97 \pm 5.89 \text{ cm}^3$ and $23.58 \pm 5.62 \text{ cm}^3$ on the right and left sides of the testis, respectively. The differences on both sides of the volumes were statistically significant. Serum hormone levels and semen parameters were significantly negatively correlated with testicular volume. There was also a statistically significant correlation between age, testicular volumes, right length, left length and left testicular width. **Conclusions:** The result from the present study shows that ultrasonography was the most accurate method for measuring the testicular volume and evaluating gonadal functions.

Keywords: Ultrasonography, Testicular volume, Anthropometric measurements, Gonadal =functions.

*Address for Correspondence:

Dr Arunkumar Neelakandan, Department of Radiology, Chettinad Hospital & Research Institute, Kelambakkam, Chennai, Tamil Nadu, INDIA.

Email: dr.arun.n@gmail.com

Received Date: 22/10/2021 Revised Date: 13/11/2021 Accepted Date: 08/12/2021

This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/). 

Access this article online

Quick Response Code:	Website: www.medpulse.in
	DOI: https://doi.org/10.26611/10132034

INTRODUCTION

The measurement of testicular volume in children, adolescents and adults are essential in the preliminary examinations for gonadal function since it correlates well

with various testicular function indices. The testes are responsible for the production of spermatozoa and the secretion of testosterone in the man. Approximately 80–90% of the testicular volume is secreted by seminiferous tubules and germ cells.^{1,2} Thus, a reduction in the number of these cells is manifested in a reduction in testicular volume. The assessment of testicular volume has been extensively studied in recent years. The testicular volume has traditionally been obtained using instruments such as the Prader or punched-out orchidometer. Currently, many measurement methods employed include callipers, orchidometry, and ultrasonography.³ Ultrasonography is the most readily available and accurate method of measuring testicular volume as determined by comparison with the actual volume. However, earlier many studies have shown that testicular volume measured by using

ultrasonography varies widely depending on the formula used.¹⁻⁵ Reliable and accurate testicular volume is greatly beneficial in evaluating patients with disorders affecting testicular growth, development, and function. Studies in infertile men have shown that the testicular volume directly correlates to seminal fluid and sex hormone assay, just like the simple measurement of testicular length, width and depth.⁴⁻⁶ A total testicular volume (i.e., summation of right and left) of 20 ml and more, as determined by ultrasound, indicates normal testicular function. In adult males, testicular volume is measured, which is directly related to the spermatogenic activity. Testicular volume is related to various reproductive endocrine parameters. Measurement of testicular size and volume is connected with the assessment of male fertility. The seminiferous tubules comprise 70% to 80% of the testicular mass. The testicular volume is believed to be an index of spermatogenesis.⁷ In contrast, testicular volume measurement in adolescent boys is vital in assessing the onset of puberty or pubertal development. It is also used to evaluate boys with various disorders affecting testicular growth and development, such as varicocele, cryptorchidism and after testicular torsion.^{7,8} A wide range of anthropometric parameters has been proposed in human subjects. They have been widely used as an index for measuring nutritional status, gonadal growth, development and functions and have been used to predict well-being and risk of some gonadal dysfunctions.⁹ Therefore, the present study attempts to critically assess the accuracy of ultrasonography in measuring the testicular volume and verify the accuracy of the Lamberts formula for calculating the ultrasound-determined testicular volume in healthy adults.

MATERIALS AND METHODS

Our study was a prospective observational category study. The institutional scientific and ethical committee approved this study. Recruitment of Subjects: This study was conducted during August 2019 and August 2021. It was carried out at Department of Radiology, Chettinad Hospital and Research Institute, Kelambakkam, Chennai.

RESULTS

In this study, the subjects' mean (\pm SD) age was 38 years, most 23 of whom belonged to the 30–39 years age group. Out of 100 subjects, 57 subjects were married, 31 were single, and 12 were widowed/divorced. A total of 47 subjects were educated up to graduation level, and only nine subjects were from the uneducated category. Twelve participants reported a positive family history of gonadal dysfunction/infertility (Table 1).

Table 1: General Demographic features of adult healthy subjects enrolled in this study.

Parameters	Subjects (n=100)	
Age (Years)	18-29	22
	30-39	23
	40-49	20
	50-59	19
	60-70	16

A total of 100 healthy adult males aged 18 to 70 years were assessed to measure the testicular size, and Lambert's formula was used to calculate testicular volumes on both sides. The relationship between testicular measurements and other participants' characteristics was also evaluated. A total of 100 healthy adult males aged 18 to 70 years (median age 38 years) were recruited for this study. Those who have willingly signed the informed consent form and self-declaration related that he does not suffer from any other metabolic disorders were considered for this study. Subjects with a previous history of scrotal surgery, genital or scrotal abnormality and those being investigated for infertility were not considered for this study. Patient position: Supine position for the ultrasonography of each subject was used. Biochemical Estimation: Serum Hormonal Analysis was performed by electrochemiluminescence with Hitachi-Roche equipment (USA). Blood sampling was performed at 8.00 am, after at least 8 hours of sleep. Determination of serum LH and prolactin was repeated at a distance of 30 minutes. Semen Analysis: Semen analysis was carried out according to the WHO criteria. The various parameters were examined which includes colour, volume, liquefaction time, pH, density, total count, progressive motility, morphology, and leukocytes. Imaging examination: All the 100 subjects were evaluated with the help of the aid of the 5.5-7.5MHZ linear transducer from a Philips HD11XE ultrasound machine [Here, you can mention the ultrasound machine details used in that particular hospital]. The sagittal length, width and transverse height were measured, all in centimetre (cm). The sagittal diameter, characterised by the mediastinum, was identified as an echogenic line running from the superior to the inferior pole of the testis. In sagittal diameter, the epididymis was projected separately from the testis. The epididymis was not included in the volume measurement. Lambert's formula⁴, $W \times H \times L \times 0.71$, was then used to calculate the volume for both the right and left testes. The measurements were recorded on the questionnaire (datasheet) for each patient on each examination day.

Statistical analysis: The consolidated and compiled data were analysed with SPSS statistics software.

Gender	Male	100
Marital Status	Married	57
	Unmarried	31
	Widowed/Divorced	12
Education Level	Illiterate	9
	Up to 10th std.	21
	Graduate	47
Socioeconomic status	Postgraduate	23
	Lower	16
	Middle	73
Occupation	Upper	11
	Student	18
	Service	48
	Self-employed	22
Residence area	Retired	12
	Rural	58
	Urban	42
Family history of Gonadal dysfunction / Infertility	Yes	12
	No	88

The variations in the testicular volume of both the right and left testis in subjects were normally distributed, as highlighted in Table 2. Mean right and left testicular volumes were $21.97 \pm 5.89 \text{ cm}^3$ and $23.58 \pm 5.62 \text{ cm}^3$, respectively. The left testis's mean length, width, and height were $3.55 \pm 0.28 \text{ cm}$, $2.49 \pm 0.33 \text{ cm}$, and $2.61 \pm 0.30 \text{ cm}$, and The right testis's mean length, width, and height were $3.53 \pm 0.30 \text{ cm}$, $2.46 \pm 0.34 \text{ cm}$, and $2.57 \pm 0.30 \text{ cm}$, respectively. Although the left testis's length, width, and volume were larger than the right, these differences were statistically significant ($p > 0.05$).

Table 2: Average of measurement of general parameters of testes among all study subjects.

Variables	Left Side	Right Side
Height (cm)	2.61 ± 0.30	2.57 ± 0.30
Width (cm)	2.49 ± 0.33	2.46 ± 0.34
Length (cm)	3.55 ± 0.28	3.53 ± 0.30
Volume (cm³)	23.58 ± 5.62	21.97 ± 5.89

Data represents Mean \pm SD values.

A comparison of testicular volumes by age, groups on both sides, is shown in Table 2. Across the age groups, the right testicular volumes were predominantly larger than the left side. However, in all the age groups, the results of a paired-samples t-test, as highlighted in the table, showed no significant differences on both right and left sides ($p > 0.05$).

Table 3: Relationship between subjects age, height, weight and body mass index (BMI)

Age (Years)	N	Height (m)	Weight (Kg)	BMI (Kg/m ²)
18-29	22	1.49 ± 0.32	63.4 ± 8.2	22.0 ± 1.7
30-39	23	1.56 ± 0.49	64.2 ± 5.7	24.9 ± 1.8
40-49	20	1.53 ± 0.51	69.4 ± 6.8	25.3 ± 2.2
50-59	19	1.64 ± 0.74	73.8 ± 8.4	25.5 ± 1.8
60-70	16	1.49 ± 0.39	59.4 ± 7.2	24.9 ± 2.6

Data represents Mean \pm SD values.

The results analysis of variance to detect any significant differences or trends in testicular volume compared with BMI and age groups of subjects are presented in Table 3. The age group of 18-29 years had a maximum number of underweight subjects, and the age group of 40-49 years and 50-59 years had the maximum number of obese subjects.

Table 4: Assessment of correlation between subjects' body mass index (BMI) categories and testicular volumes.

BMI category	N	Left testicular volume (cm ³)	Right testicular volume (cm ³)
Underweight	5	21.47 ± 4.12	21.07 ± 3.21
Normal	57	22.54 ± 5.27	22.48 ± 4.79
Overweight	26	23.12 ± 3.23	22.97 ± 5.29
Obese	12	23.95 ± 2.98	23.16 ± 3.34

The results of correlation analysis to evaluate; further, the relationship between BMI and testicular volume measurements are presented in Table 4. Although there appears to be a gradual increase in left testicular volume from $21.47 \pm 4.12 \text{ cm}^3$ in the underweight category to $23.95 \pm 2.98 \text{ cm}^3$ in the obese, this trend was not statistically significant ($p > 0.05$). Similarly, the right testicular volume increased from $21.07 \pm 3.21 \text{ cm}^3$ to $23.16 \pm 3.34 \text{ cm}^3$, it was also not statistically significant ($p > 0.05$). There is generally a positive but weak correlation between BMI and testicular length, width, height, and volume on both sides. These showed that as the BMI increased, testicular dimensions increased.

Table 5: Analysis of various semen parameters from the participant's group.

Parameter	Subjects (n=100)
Sperm concentration (million/mL)	43.1 ± 7.2
pH	7.2 ± 0.9
Progressive motility (%)	37.1 ± 6.2
Normal forms (%)	12.8 ± 1.9
Immature germ cells (%)	3.8 ± 0.4
Volume (mL)	2.9 ± 0.7

The conventional sperm parameters of the participants enrolled in this study were shown in Table 5. These patients with low testicular volume had a statistically significant lower seminal fluid volume, sperm concentration, and percentage of progressive motility and spermatozoa with normal form.

Table 6: Analysis of various serum hormone levels among the participant's group.

Parameter	Subjects (n=100)
LH (mIU/mL ⁻¹)	5.1 ± 0.7
FSH (mIU/mL ⁻¹)	4.6 ± 0.8
Total testosterone (ng/mL ⁻¹)	5.9 ± 0.9
Estradiol (pg/mL ⁻¹)	26.2 ± 3.9
Prolactin (ng/mL ⁻¹)	17.1 ± 3.1

The hormonal serum levels are displayed in Table 6. LH, FSH, and prolactin levels were similar among all the participants.

DISCUSSION

The measurement of testicular volume in adults is an essential parameter in evaluating the health status of the testes in various clinical conditions such as undescended testis, torsion, malignancies, orchitis and varicocele. In addition, the estimation of testicular volume is an integral aspect of male infertility evaluation.¹⁻³ Compared to Prader orchidometry or the punched-out orchidometer, ultrasound has generally been recognized to be the most accurate^{10, 12, 39} and testicular volume measurements using the ultrasonographic formula $L \times W \times D \times 0.71$, which was employed in this study, has been reported to be the closest to actual testicular volumes in humans.¹⁻⁵ Moreover, the non-invasiveness of ultrasound and the absence of concerns about radiation allow for repeated evaluations. The sample selection criteria used in this study are similar to those used in earlier studies.⁶⁻⁹ In these studies and the current one, participants with a history of testicular and scrotal surgery or disease were excluded from the study. The mean right and left testicular volumes obtained in this study were $21.97 \pm 5.89 \text{ cm}^3$ and $23.58 \pm 5.62 \text{ cm}^3$, respectively. This study also revealed an increasing trend in volumes as age advances to a peak in the late 30s, followed by a decrease in the volumes. This observed peak testicular volume at this period of life is consistent with 57 reports that have suggested that men are at their peak fertility at this period and that beyond 50 years of age,

serum testosterone and spermatogenesis reduce with time.¹⁰ More detailed analysis showed no statistically significant differences in these trends for both right and left testes. Age was also positively but weakly correlated to the right length, left length and left testicular width. A negative and weak correlation was obtained between age and left testicular volume. This is different from the results obtained in other studies,⁷⁻¹⁰ where age showed no correlation with the volume of the left and the right side of the testis. Few earlier studies had established that the testes achieved their maximal size by 18 years and remained so until 80 years when they started decreasing in size.⁸⁻¹² Further, the correlation analysis revealed a positive but weak correlation between BMI and testicular length, width, height and volume,¹³⁻¹⁵ establishing a weak and positive correlation between testicular volume and body mass index. However, few studies.⁹⁻¹³ showed a consistent lack of correlation between BMI and left testicular volume. Furthermore, as argued by Sotos and Tokar,⁷ the measurement of the testicular volume is not an exact science. Even with the reliability of the ultrasound, variability related to the transducer used, the possibility of compression of the testis, and intra- and inter-observer variation in the measurements (width, height, length, and volumes), among other factors, exists. In addition, other reasons such as environmental, nutritional and genetic factors may be responsible for these differences.¹⁴⁻¹⁶ In

general, in this study, it was found that the testicular volume had a positive but weak correlation with weight, height, BMI, but none was statistically significant. Although the left testis's length, width, and volume were larger than the right, these differences were statistically significant. However, no immediate scientific explanation could be inferred to the differential association of the right and left testes with the measured anthropometric parameters. This variation warrants further studies in this subject area to establish a clinical and scientific basis of the variations in right and left side testicular volumes. The results of this study indicate that participants with a reduced testicular volume, in the absence of testicular disease, show poorer semen parameters. The participants enrolled in this study did not show any frank endocrine alteration, but total testosterone serum concentration was lower in men with reduced testicular volume.

CONCLUSION

In this study, the right testicular volume showed less volume compared to the left testicular volume. Both side measurements have shown a positive but weak correlation with weight, height, BMI, but none was statistically significant in healthy adults. Further studies are needed to support or negate these findings. This anthropological study addressed an oft-repeated question about the accuracy of the different non-invasive methods of testicular measurements. A rough estimate of testicular size may suffice in cases that do not need any active intervention. Further studies are needed to clarify the relationship between reduced testicular volume and ultrastructural alterations of the testicular parenchyma in patients with low testicular volume and without evidence of disease at the clinical and ultrasound examinations. This study complies with the published data so far and revealed ultrasonography to be the most accurate and objective in vitro method of assessing testicular volume.

REFERENCES

- Oehme NH, Roelants M, Bruserud IS, et al. Low BMI, but not high BMI, influences the timing of puberty in boys. *Andrology*. 2021;9:837–845.
- Bellurkar A, Patwardhan S, Patil B, Kanbur A, Jain H, Velhal R. Role of testicular size as a parameter for predicting infertility in Indian males. *J Hum Reprod Sci* 2020;13:114-6.
- Innocent MC, Asomugha LA, Ukamaka MN, Aronu ME. Ultrasound measured testicular volume in Nigerian adults: Relationship of the three formulae with height, body weight, body-surface area, and body-mass index. *Int J Adv Med Health Res* 2016;3:85-90.
- Abu, S., Kolade-Yunusa, H. O., Atim, T., Obakeye, F. E., and Dakum, N. K. Correlation Between Ultrasonic Testicular Volume and Seminal Fluid Analysis in Men with Infertility. *European Journal of Medical and Health Sciences*, 2021;3(1), 103–107. <https://doi.org/10.24018/ejmed.2021.3.1.541>.
- Liu C, Liu X, Zhang X, Yang B, Huang L, Wang H and Yu H. Referential Values of Testicular Volume Measured by Ultrasonography in Normal Children and Adolescents: Z-Score Establishment. *Front. Pediatr.*2021;9:648711. doi: 10.3389/fped.2021.648711.
- Anyanwu, L.J.C., Sowande, O.A., Asaleye, C.M. et al. Testicular volume: correlation of ultrasonography, orchidometer and caliper measurements in children. *Afr J Urol* 2020; 26:6. <https://doi.org/10.1186/s12301-020-0016-z>.
- Sotos, J.F., Tokar, N.J. Appraisal of testicular volumes: volumes matching ultrasound values referenced to stages of genital development. *Int J Pediatr Endocrinol* 2017; 7. <https://doi.org/10.1186/s13633-017-0046-x>.
- Elder CJ, Langley J, Stanton A, De Silva S, Akbarian-Tefaghi L, Wales JKH, Wright NP. A simulation study assessing the accuracy and reliability of orchidometer estimation of testicular volume. *Clin Endocrinol (Oxf)*. 2019 Apr; 90(4):623-629. doi: 10.1111/cen.13923.
- Nassan FL, Jensen TK, Priskorn L, Halldorsson TI, Chavarro JE, Jorgensen N. Association of Dietary Patterns With Testicular Function in Young Danish Men. *JAMA Netw Open*. 2020 Feb 5; 3(2):e1921610. doi: 10.1001/jamanetworkopen.2019.21610.
- Sakamoto H, Saito K, Oohta M, Inoue K, Ogawa Y, Yoshida H. Testicular volume measurement: comparison of ultrasonography, orchidometry, and water displacement. *Urology*. 2007 Jan; 69(1):152-7. doi: 10.1016/j.urology.2006.09.012.
- K.H. Tijani, B.O. Oyende, G.O. Awosanya, R.W. Ojewola, A.O. Yusuf. Assessment of testicular volume: A comparison of fertile and sub-fertile West African men. *African Journal of Urology* 2014; 20(3), 136-140.
- Boehme NHB, Roelants M, Saervold Bruserud I, et al. Reference data for testicular volume measured with ultrasound and pubic hair in Norwegian boys are comparable with Northern European populations. *Acta Paediatr*. 2020; 109:1612–1619. <https://doi.org/10.1111/apa.1515>.
- Tseng, CS., Huang, KH., Kuo, MC. et al. The impact of primary location and age at orchiopexy on testicular atrophy for congenital undescended testis. *Sci Rep* 2019; 9:9489. <https://doi.org/10.1038/s41598-019-45921-6>.
- Manuel, B., W. Ugboma, E., and C. Nwankwo, N. Relationship between Testicular Volume and Sperm Count in Infertile Men in Southern Nigeria. *Asian Journal of Medicine and Health*, 2017; 4(1), 1-6. <https://doi.org/10.9734/AJMAH/2017/29182>.
- Boben GE, Umopathy P, Ravichandran L, Godfrey DA, Ramani G, Srinivasan V. Evaluation of testicular volume in children aged 8-17 years in south India. *Indian J Child Health*. 2016; 3(3):208-211.
- Jaiswal VK, Khadilkar V, Khadilkar A, Lohiya N. Stretched penile length and testicular size from birth to 18 years in boys from Western Maharashtra. *Indian J Endocr Metab* 2019; 23:3-8.

Source of Support: None Declared
Conflict of Interest: None Declared